

Treatment for insertional Achilles tendinopathy: a systematic review

J. I. Wiegerinck · G. M. Kerkhoffs ·
M. N. van Sterkenburg · I. N. Sierevelt ·
C. N. van Dijk

Received: 28 February 2012 / Accepted: 17 September 2012 / Published online: 6 October 2012
© Springer-Verlag Berlin Heidelberg 2012

Abstract

Purpose Systematically search and analyse the results of surgical and non-surgical treatments for insertional Achilles tendinopathy.

Methods A structured systematic review of the literature was performed to identify surgical and non-surgical therapeutic studies reporting on ten or more adults with insertional Achilles tendinopathy. MEDLINE, CINAHL, EMBASE (Classic) and the Cochrane database of controlled trials (1945–March 2011) were searched. The Coleman methodology score was used to assess the quality of included articles, and these were analysed with an emphasis on change in pain score, patient satisfaction and complication rate.

Results Of 451 reviewed abstracts, 14 trials met our inclusion criteria evaluating 452 procedures in 433 patients. Five surgical techniques were evaluated; all had a good patient satisfaction (avg. 89 %). The complication ratio differed substantially between techniques. Two studies analysed injections showing significant decrease in visual analogue scale (VAS). Eccentric exercises showed a significant decrease in VAS, but a large group of patients was unsatisfied. Extracorporeal shockwave therapy (ESWT) was superior to both wait-and-see and an eccentric training regime. One study evaluated laser CO₂, TECAR and cryoultrasound, all with significant decrease in VAS.

Conclusions Despite differences in outcome and complication ratio, the patient satisfaction is high in all surgical

studies. It is not possible to draw conclusions regarding the best surgical treatment for insertional Achilles tendinopathy. ESWT seems effective in patients with non-calcified insertional Achilles tendinopathy. Although both eccentric exercises resulted in a decrease in VAS score, full range of motion eccentric exercises shows a low patient satisfaction compared to floor level exercises and other conservative treatment modalities.

Level of evidence III.

Keywords Achilles · Insertion · Tendinopathy · Treatment · Systematic review

Introduction

Much has been published on the treatment for (midportion/non-insertional) Achilles tendinopathy and retrocalcaneal bursitis [11, 16, 23, 27, 29, 38, 39, 43, 46]. There is a tendency to prescribe known effective treatment for (midportion/non-insertional) Achilles tendinopathy or retrocalcaneal bursitis to patients with insertional Achilles tendinopathy [3, 17, 22, 28, 34, 36, 52]. Insertional Achilles tendinopathy however is a distinct clinical entity from midportion/non-insertional Achilles tendinopathy or retrocalcaneal bursitis [8, 21, 35, 45, 50, 52]. Aetiology, injury mechanism, treatment and rehabilitation differ between these pathologies [34, 48]. The ‘extrapolation’ is in part due to the confusing terminology of Achilles tendon pathology [7, 13, 25]. Different pathologies have been compared under the name of insertion tendinosis/pathy of the Achilles tendon [2, 17, 40]. Moreover, the confusing terminology aids in the current complicated overview of data for the treatment for insertional Achilles tendinopathy [2, 30, 34, 38, 41]. A recent terminology on Achilles

J. I. Wiegerinck (✉) · G. M. Kerkhoffs ·
M. N. van Sterkenburg · I. N. Sierevelt · C. N. van Dijk
Department of Orthopaedic Surgery, Academic Medical Center,
University of Amsterdam, PO Box 22700, 1100 DE Amsterdam,
The Netherlands
e-mail: j.i.wiegerinck@amc.uva.nl

tendon pathologies was proposed to simplify the current terminology and hereby aid the clinical practice, both in diagnostics and treatment [50]. The current treatment for insertional Achilles tendinopathy varies substantially; several surgical and non-surgical treatments have been described, but there is no consensus on the most effective management [5, 6, 10, 24, 42]. One previous study concluded there is a consensus that non-surgical intervention is preferred prior to surgical intervention for insertional Achilles tendinopathy [24]. However, it has not been studied which treatment is the most effective *non-surgical* intervention or which is the most effective *surgical* intervention. The purpose of this systematic review is to analyse the effectiveness of different available surgical and/or non-surgical treatment modalities for insertional Achilles tendinopathy.

Materials and methods

The definition of insertional Achilles tendinopathy, retrocalcaneal bursitis and midportion Achilles tendinopathy for this systematic review was based on the recently proposed terminology on AT pathology [50] (Table 1).

Primary outcome measures

- Change of numeric pain score [e.g. visual analogue scale (VAS)]
- Patient satisfaction

Secondary outcome measures

- Complication rate
- Coleman methodology score
- Intervention
- Diagnostic method
- Other outcome data

Inclusion criteria

Studies reporting on the outcome of treatment for insertional Achilles tendinopathy were eligible for this systematic review. A pain scale, overall patient satisfaction or American orthopaedic foot and ankle (AOFAS)-Hindfoot scale score had to be reported. Review and expert opinion papers were excluded. There were no language barriers.

Exclusion criteria

Studies on the treatment for Achilles midportion tendinopathy or unspecified Achilles tendinopathies were excluded. Studies combining the results of insertional and non-insertional Achilles tendinopathy were also excluded if the results of insertional Achilles tendinopathy could not be analysed separately. All studies on pathology located at or around the Achilles tendon insertion but not diagnosed as insertional Achilles tendinopathy (e.g. retrocalcaneal bursitis) were excluded. Studies were excluded if scaled data of pain (e.g.

Table 1 Terminology of Achilles tendon pathology

Term	Anatomic location	Symptoms	Clinical findings	Histopathology
Insertional Achilles tendinopathy	Insertion of AT onto calcaneus, most often with formation of bone spurs and calcifications in tendon proper at insertion site	Pain, stiffness, sometimes a (solid) swelling	Painful tendon insertion at the <i>midportion</i> of the <i>posterior</i> aspect of the calcaneus, swelling may be visible and a bony spur may be palpable	Ossification of enthesial fibrocartilage, and sometimes small tendon tears occurring at tendon–bone junction
Retrocalcaneal bursitis	Bursa in the recess between the anterior inferior side of the AT and the <i>posterosuperior</i> aspect of the calcaneus (retrocalcaneal recess)	Painful swelling superior to calcaneus	Painful soft tissue swelling, medial and lateral to the AT at the level of the <i>posterosuperior</i> calcaneus	Fibro-cartilaginous bursal walls show degeneration and/or calcification, with hypertrophy of the synovial infoldings and accumulation of fluid in the bursa. Alternatively, the bursa may be primarily involved by inflammatory or infectious bursitis due to an inflammatory arthropathy
Midportion Achilles tendinopathy	2–7 cm from the insertion onto the calcaneus	A combination of pain, swelling and impaired performance	Diffuse or localized swelling	Includes, but is not limited to, the histopathological diagnosis of tendinosis: implies histopathological diagnosis of tendon degeneration without clinical or histological signs of intratendinous inflammation, not necessarily symptomatic




Fig. 1 Search strategy

VAS) or patient satisfaction was not provided. Studies reporting on <10 patients were excluded.

Study population

Adults (>18 years) with diagnosed insertional Achilles tendinopathy were eligible. Adults with non-insertional or non-specified problems of the Achilles tendon insertion were excluded.

Search strategy

An extensive MEDLINE, EMBASE (Classic) and CINAHL search (1945–March 2011) was performed using the following terms: (((((insertion OR insertional OR enthesi OR attachment) AND (enthesitis OR enthesopathy OR tendinopathy OR tendinosis OR tendinoses OR tendinitis OR tendinitides OR tendinopathies OR tendonitis OR tendontides OR tenosynovitis) AND achilles)) OR ((tendon OR tendons OR tendo) AND (enthesi OR attachment OR insertion OR insertional) AND achilles AND (spur OR spurs OR calcification OR calcif*))))). In addition, the Cochrane Database of Clinical and Randomized Controlled Trials were searched using the search term 'Achilles'. After combining the results and removing duplicates of these searches, a total of 451 papers were retrieved. Based on title, abstract and study design, 393 articles were excluded. Exclusion was based on study design (105 articles), papers unrelated to the Achilles tendon (172), related to the Achilles tendon but unrelated to insertional Achilles

tendinopathy (33 articles related to midportion tendinopathy, 83 articles related to Achilles tendon rupture) (Fig. 1). The full text of the remaining 58 articles was obtained. A review of references of full text articles obtained 15 new articles. A total of 73 full text articles were reviewed. 59 articles were excluded based on full text, resulting in the inclusion of 14 articles for this review (Table 3) [10, 12–14, 20, 22, 26, 31, 32, 37, 42, 44, 51, 53].

Data collection and study selection

Two reviewers independently assessed all studies. These two reviewers identified all titles, assessed the abstracts independently and excluded the irrelevant articles. Any remaining difference between reviewers was settled by discussion. Authors of the included studies were contacted by e-mail or phone to retrieve any necessary additional data (Tables 3, 5).

Data extraction

The two reviewers independently extracted data for the aforementioned outcome using a data extraction form. Differences were settled by discussion. Complications were divided in minor and major (Table 2).

Statistical analysis and quality assessment

Data are presented as weighted means and summed percentages. In addition, all included studies were measured

Table 2 Classification of complications

Major complications	Minor complications
Achilles tendon rupture	Discomfort
Any reoperation	Superficial infections
Deep venous thrombosis	Minor wound problems
Reflex dystrophy	Scar tenderness/hypertrophy
Persisting neuralgia	Mild form of paresthesia
Deep infections/major wound problems	Prolonged hospitalization

according the Coleman methodology score (CMS) for tendinopathies of the Achilles tendon (Appendix 1) [47]. This score was created to assess the quality of studies evaluating the surgical outcome for Achilles tendinopathies [47]. The scale consists of two parts, with a highest possible score of 100. To simplify the quality comparison of surgical and non-surgical studies, the ‘description of surgical procedure given’ was interpreted as ‘description of procedure given’ in non-surgical studies. In addition, as there is no post-operative rehabilitation in non-surgical studies, a maximum score was granted to non-surgical studies for “description of post-operative rehabilitation”. The authors of the score stated that an increase in CMS would show a decrease in effectiveness of the evaluated treatment [47]. It was also described that an increase in CMS had a positive linear correlation with an increase in year of publication [47]. The Spearman correlation coefficient (ρ) is calculated for both hypotheses.

Results

Results are divided in a surgical and a non-surgical group. The non-surgical group is subdivided in four groups.

Population characteristics

The total number of patients with insertional Achilles tendinopathy in the 14 included studies was 433; 452 tendons were treated. Overall, the weighted mean age of 12 studies was 49 years (range 18–83), 51 % male and 49 % female (two studies did not report patient characteristics). The subgroup surgical treatment had a weighted mean age of 54 (range 18–83) years (one study did not report on patient characteristics). The non-surgically treated subgroup consisted of 50 % male and 50 % female with a weighted mean age of 44 (range 21–80) years (one study did not report on patient characteristics).

Surgical treatment

Six studies, covering five operative techniques and a total of 200 patients with 211 tendons (procedures), were

evaluated (Tables 3, 4) [12, 20, 31, 33, 51, 53]. The weighted mean age of surgically treated patients was 54 (range 47–57). The male–female ratio was 6:5 (55:45 %). Five studies reported on patient satisfaction, all showed a relative good outcome regarding patient satisfaction with 163 out of 183 patients (89 %) reporting an excellent or good satisfaction (Table 4) [12, 31, 32, 51, 53]. A total of 52 complications (23 %) were documented; including seven major complications (3.1 %) (Table 4). Subjective pain scales and other questionnaires were not regularly used (Tables 3, 5).

Non-surgical treatment

Injections

Two studies evaluated different types of injections [37, 44]. In a retrospective analysis, Ryan et al. [44] included 22 patients treated with ultrasound (US)-guided hyperosmolar dextrose (20 mg/mL) injections. A mean of 5 injections were given. The mean VAS on daily activity decreased with 4.1 points ($p = 0.001$) (on a 10-point VAS scale) at follow-up (28.6 months). The authors did not report on patient satisfaction. Ohberg et al. [37] prospectively analysed 11 patients, who were treated with polidocanol injections until symptoms dissolved (up to five injections). The mean VAS showed a decrease of 5.9 at 8 months follow-up, and 8 out of 11 (73 %) patients were satisfied with the outcome (Table 5).

Eccentric training

Four studies evaluated the treatment for eccentric exercises [13, 22, 26, 42]. A total of 92 patients with 100 painful tendons were studied. A 12-week daily eccentric treatment regime was used in every study. There were some differences in treatment protocol; three studies evaluated full range eccentric exercises (below the step), whereas the other evaluated floor level eccentric exercises (providing less stress on the Achilles’ insertion); in addition, there were some difference between ‘number of daily sets’ and ‘number of repetitions’. All four studies used a numerical pain scale; three studies documented the patient satisfaction of the received treatment [13, 22, 42]. When combining the results, an overall decline on a 10-point scale was 2.7 points (weighted mean). All studies combined, and the patient satisfaction showed a large group of unsatisfied patients: of 83 patients only 35 (42 %) were either ‘extremely satisfied’ or ‘satisfied’ (Table 5). Subdividing the full range of motion eccentric group from the floor level eccentric exercises, a noticeable difference was found. The VAS for full range of motion decreased with a weighted mean of 2.0 points, whereas in the floor level, group is

Table 3 Overview of included studies

Author, year	Design (level of evidence)	N patients (N tendons)	Age (range) in years	Method of diagnosis	Duration of symptoms	Follow-up in month (range)	Bony disorder?	Intervention 1	Intervention 2 or more	Outcomes
Costantino et al. [10]	III	15 (15)	32.7*	US*	>2 month	8 month	NR*	Lacer CO ²	Int. 2: TECAR Int. 3: Cryoultrasound	VAS; patient satisfaction
Elias et al. [12]	IV	40 (40)	57 (39–76)	MRI/US	>6 month	27 month (18–68)	“Haglund” exostosis	Surgical (Table 3)	–	VAS; AOFAS; patient satisfaction
Fahlstrom et al. [13]	III	30 (31)	37.9 (SD ± 11.6)	Varies	32 month (42.4 SD)	3 month	Mixed pattern	12 week eccentric training: 7d/w 2p/d	–	VAS; patient satisfaction
Furia [14]	III	68 (68)	50 (21–80)	X-lat	19.9 month (6–60)	12 month	Supr/ calcification on X-ray	1 × ESWT (3,000 shocks: 604 mJ/mm ²)	Matched control group n = 33	VAS; roles and maudslsey score
Johnson et al. [20]	IV	22 (22)	51 (45–75)*	X-lat*	36 month (2–216 month)	34 month (11–64)	Calcification at AT insertion*	Surgical (Table 3)	–	AOFAS
Jonsson et al. [22]	III	27 (34)	53.4 (25–77)	US	26.5 month (6–96)	4 month	Calcification/spur	12 week painful ecc. Training: 7d/w 2p/d	–	VAS; patient satisfaction
Knobloch [26]	III	10 (10)	NR*	US	NR*	3 month	No/excl	12 week painful ecc. training: 7d/w 3p/d (15 reps)	–	VAS
Maffulli et al. [31]	III	21 (21)	46.9 (38–59)	X-lat + US	20.2 month (14–45)	48.4 month (24–48)	Calcific ins. Tend.	Surgical (Table 3)	–	VISA-A; patient satisfaction
McGarvey et al. [32]	IV	21 (22)	54.4 (30–77)	Varies*	>6 month*	33 month (26–40)	Mixed pattern*	Surgical (Table 3)	–	Pain (no scale); patient satisfaction
Ohberg and Alfredson [37]	III	11 (11)	44 (24–67)	US and colour doppler	29 month (7–96)	8 month	Some w. bony enlargement	Sclerosing polidocanol (5 mg/mL) injection	–	VAS; patient satisfaction
Rompe et al. [42]	I	50 (50)	Int 1: 39.2 (SD ± 10.7) Int2: 40.4 (SD ± 11.3)	US	6 month (8.3–26.3)	4 month	Excl	12 week painful ecc. training: 7d/w 3p/d (15 reps)	Shockwave therapy. 3 sessions of 2.5 bar	VISA-A; patient satisfaction
Ryan et al. [44]	III	22 (22)	52.7 (SD ± 11.5)*	US	21 month (7–228)	28.6 month (12–48)	NR*	US guided hypertonic dextrose (20 mg/mL) injection	–	VAS
Wagner et al. [51]	IV	61 (65)	52.7	X-lat	6 weeks	Int 1: 47 month Int 2: 33 month	NR*	Surgical (Table 3)	Surgical (Table 3)	“Complete pain relieve”; patient satisfaction
Yodowski et al. [53]	IV	35 (41)	54 (18–83)	X-lat*	6 month	36 month	“Haglund” exostosis*	Surgical (Table 3)	–	7 point pain scale; patient satisfaction

Outlined are level of evidence; patient demographics; method of diagnosis, study follow-up, reported presence of bony disorder, studied intervention and used outcome measures

NR not reported, X-lat conventional lateral radiograph of the ankle, US ultrasonography

* Authors were contacted to provide these data

Table 4 Studies describing surgical techniques, outlined are number of procedures; major and minor complications and surgical technique

Author	N of procedures	Complications		Surgical technique		
		Major	Minor	Positioning	Approach	Description of procedure
Elias et al. [12]	40	0	2	Prone	Longit. midline incision over AT	Diseased part of AT resected. Post sup. calcaneal osteotomy + spur resection. If 50–100 % of AT detached reattachment and FHL autograft
McGarvey et al. [32]	22	0	9	Prone	Longit. midline inc. through AT and insertion	Calcification debridement if excessive (>50 %) plantaris augm. or bone anchors if necessary; retrocalcaneal bursectomy, AT defect approximated with absor. sutures
Johnson et al. [20]*	22	1 [†]	2	Same technique as McGarvey		
Maffulli et al. [31]	21	0	6	Prone	Medial longit. inc. up to 1/3 of AT	Bursectomy, abnormal AT appearance were explored by tenotomise (not repaired) and excised, if 1/3 or > of AT was detached reattached with bone anchors
Wagner et al. [51]**	47	2 [±]	2	Prone	Medial J incision	AT and paratenon debridement, no/partial (<50 %) AT detachment; sup calcaneal tuberosity resection. Reattachment with 4 bone anchors
Wagner et al. [51]**	33	3 [§]	7	Prone	Medial J incision	AT and paratenon debridement, complete AT release; sup calc. tuberosity resection; modified Krakow technique, VY lengthening reattachment with 4 bone anchors
Yodlowski et al. [53]	41	1 [‡]	17	Prone	Lateral longit. Inc. between AT and supero- lateral crest of calcaneus	Preop AB iv. Fat pad; bursa; sup calc. tuberosity resection. Excision of scar tissue; adhesions between AT and paratenon. Check for intratendinous calcifications, excised if present. Reattachment of AT with bone anchors if necessary
Totals	226	7 (3.1 %)	45 (20 %)			

AT Achilles tendon, AB antibiotics

* Same surgical technique as McGarvey

** Same study comparing complete detachment of the AT with partial or no detachment

± One substantial wound infection; one surgical revision (complete detachment)

§ One AT rupture; two major wound problems

† One deep venous thrombosis

‡ One deep venous thrombosis

decreased with 3.9 points. In addition, 18 of 27 (67 %) were satisfied in the floor level group, in the full range of motion only 17 of 56 (30 %) were satisfied (Table 5).

Extracorporeal shockwave therapy

Two studies tested the effect of extracorporeal shockwave therapy (ESWT) for insertional Achilles tendinopathy, one prospective study and one randomized controlled trial [14, 42]. Furia et al. [14] prospectively compared an ESWT group to a matched control group receiving non-specified 'traditional treatment'. A mean decrease in VAS of four points was measured after 12 months; 29 out of 35 (83 %) patients were satisfied with ESWT. The control group ($n = 33$) showed a significantly less decrease in VAS and

patient satisfaction. Rompe et al. [42] compared ESWT with an eccentric training regime in a RCT, 25 patients received ESWT. The VAS decrease was 5.1 points at final follow-up, 16 (64 %) patients were satisfied with their treatment. Overall, 49 out of 64 (77 %) were satisfied after ESWT (Table 5).

Others

One study evaluated three different non-surgical treatment modalities: Costantino et al. [10] compared laser CO² with TECAR and cryoultrasound. Five patients were included in each group. Every intervention showed a significant decrease in VAS score at follow-up. All five patients in the cryoultrasound group were very satisfied; in the TECAR

Table 5 Results of different treatment modalities for insertional Achilles tendinopathy (IAT)

	N Patients (tendons)	Age in yrs	F-U (months)	VAS decrease at F-U	AOFAS increase at F-U	VISA-A increase at F-U	Patient satisfaction		CMS
							Excel/good	Fair/poor	
Surgical interventions for IAT									
Elias et al. [12]	40 (40)	57	27	-7.2	39.9	-	38	2	65
Johnson et al. [20]	22 (22)	51*	34	-	36	-	-*	-*	48
Maffulli et al. [31]	21 (21)	46.9	48.8	-	-	22.6	16	5	67
McGarvey et al. [32]	21 (22)	54.4	33	13 no pain; 4 less pain; 3 unchanged; 2 worse	18	4	36	-	-
Wagner et al. [51]*	22 (26)	55	47	-	-	-	24	2	52
Wagner et al. [51]*	39 (39)	56	33	-	-	-	36	3	-
Yodkowski et al. [53]	35 (41)	54	36	7 pnt pain scale: pre-op: 4.7 post-op: 1.5	31	4	46	-	-
Totals	200 (211)						163 (89 %)	20 (11 %)	Mean: 52
Weighted Mean	54		35.6		38.5				
Injections for IAT									
Ohberg and Alfredson [37]	11 (11)	44.0	8	-5.85	-	-	8	3	55
Ryan et al. [44]	22 (22)	52.7*	4	-4.1	-	-	-*	-*	65
Totals	33 (33)						8 (73 %)	3 (27 %)	Mean: 60
Weighted Mean	44		5.3	-4.7	-	-			
Eccentric training for IAT									
Fahlstrom [13]	30 (31)	37.9	3	-2	-	-	10	21	59
Jonsson [22]	27 (34)	53.4	4	-3.9	-	-	18	9	63
Knobloch [26]	10 (10)	-*	3	-2.8	-	-	-*	-*	65
Rompe et al. [42]	25 (25)	39.2	4	-1.8	-	10	7	18	78
Totals	92 (100)						35 (42 %)	48 (58 %)	Mean: 66
Weighted mean	44.1		3.6	-2.67	-	-			
Extracorporeal shock wave therapy for IAT									
Furia [14]	35 (35)†	50.0	12	-4	-	-	29	6	64
Rompe et al. [43]‡	25 (35)	40.4	4	-5.1	-	27	16	9	78
Totals	60 (60)						45 (75 %)	15 (25 %)	Mean: 71
Weighted mean	46		8.7	-4.64	-	-			
Other non-surgical treatment for IAT									
Costantino	15 (15)	32.7	8	-6.2	-7.0	-7.2	-	-	15* (100 %)
Total of all	400 (419)						255 (75 %)	86 (25 %)	Mean: 58
Weighted mean	49		21.2						

Studies are ordered by treatment modality, with respective subtotals. The overall total is presented at the bottom

F-U follow-up, CMS Coleman methodology score

- Data not available

† Matched control group (n = 33) is not included in this table

‡ Same study

* Authors were contacted to provide this data




Fig. 2 Relationship between CMS and reported success rate (non-significant difference)

group, two were very satisfied and three were satisfied; in the laser CO² group, all five were satisfied (Table 5).

Methodological quality

All included studies were measured according to the CMS for tendinopathies of the AT [47]. The mean CMS was 58 (range 36–78) (Table 4). The surgical studies scored a mean CMS of 52 (range 36–67), all non-surgical studies scored a mean CMS of 62 (range 44–78). We did not find a significant ($p < 0.05$) correlation between the CMS and effectiveness of the evaluated treatment (Fig. 2). In addition, the authors of the CMS stated there is a positive linear correlation between the date of publications and the CMS [47]. We found a significant positive linear correlation of 0.71 ($p = 0.02$) (Fig. 3).

Discussion

The most important findings of this study are the outcome differences between ESWT, full range of motion eccentric exercises, floor level eccentric exercises, and furthermore the overall good outcome after surgical treatment in case of persisting complaints. One previous systematic review was performed regarding insertional Achilles tendinopathy, this included 11 articles and outlined the current used treatments; however, no outcome data were provided [24]. Despite the overlap, we believe that this systematic review adds methodological quality and incorporates recently published data to provide the best currently available evidence. Regardless of the strict inclusion criteria, our systematic review reports on 14 articles and specifically emphasizes the patient satisfaction and change in pain scores. Herewith, we aimed to provide an overview of the effectiveness for every available treatment for insertional Achilles tendinopathy. Kearney et al. [24] concluded that




Fig. 3 Relationship between CMS and year of publication $\rho = 0.71$ (significant difference, $p = 0.02$)

non-surgical treatment is preferred prior to surgical treatment. In addition, they concluded that non-surgical treatment favoured eccentric training and ESWT [24]. The results of the current review however showed a large group of unsatisfied patients after eccentric training compared to other non-surgical treatment [12, 13, 22, 42] (Table 4). The reason of this difference in interpretation of almost equal data is unsure. Kearney et al. did not include multiple studies which were included in this review. In addition, their (primary) outcome measures were not clearly stated. The primary outcomes of the current study are patient satisfaction and change in pain scale. Hence, the primary outcomes variance may cause the interpretation differences. As eccentric training has a substantial effect on the decrease of pain, this may be interpreted as effective; Kearney et al. may have based their conclusion hereupon regardless of the patient satisfaction. The included studies used a wide range of outcomes with little consistency between studies (Tables 3, 5). Evaluation and comparison are therefore difficult. As data are presented as weighted means and summed percentages, the results should be interpreted with caution. The most frequent used outcome was a numerical pain scale, often a 11-point scale but also a 7-point or no numerical scale was used. In addition, the AOFAS, VISA-A, Roles & Maudsley score and different patient satisfaction measurement methods were used. The CMS was used to assess the methodological quality of included articles. The CMS was first designed to assess the quality of patella tendinopathy, later it was modified by Tallon et al. to assess the quality of surgical studies on Achilles tendinopathy [9, 47]. Out of a maximum score of 100 point, the mean CMS for included studies was 58 (range 36–78). As stated, the maximum score of ten points was granted for ‘description of post-operative rehabilitation’ to non-surgical studies. This simplified the comparison of relative outcome in quality between surgical and non-surgical studies. It also implicates that the absolute

mean overall score of non-surgical studies would be biased by the used treatment as these studies start with a CMS of ten points. All included surgical studies however scored ten points for ‘description of post-operative rehabilitation’.

Surgical treatment

A total of six studies evaluated the outcome of surgical treatment for insertional Achilles tendinopathy [12, 20, 31, 32, 51, 53]. The evaluated studies described five different operative techniques (Table 4). None of the included surgical techniques were minimally invasive. All minimal invasive surgical studies were excluded due to unclear terminology, minimal invasive studies focused on retrocalcaneal bursitis instead of insertional Achilles tendinopathy [29, 49]. Three studies used a medial incision: two a midline or central incision and one a lateral incision. All techniques involved resection of the posterosuperior calcaneal prominence, retrocalcaneal bursa and intratendinous calcifications. All studies detached the AT partially or in total, reinserted the AT with bone anchors or performed augmentation (plantaris tendon or FHL). Despite the (minor) differences in surgical technique, there were large differences in complication rate. One study reported no major and only 7.5 % minor complications [12]. Others reported 23 % major and 59 % overall complications [32]. Although the difference in complication rate was evident, there was almost no difference in patient satisfaction (Table 5). Outcome on pain scores (e.g. VAS) was scarcely reported [12, 53]. Based on the included studies, it is not possible to draw any conclusions regarding the best surgical technique for insertional Achilles tendinopathy.

Non-surgical treatment

Injections

Two studies evaluated medicinal injections as a minimal invasive treatment [37, 44]. Sclerosing polidocanol was injected as well as hyperosmolar dextrose. As the included study with sclerosing polidocanol is only a pilot, with hitherto no subsequent study, the positive outcome should be considered with care. Hyperosmolar dextrose injections for tendinopathies have been studied rarely, albeit the positive results raise expectations, clinical use should be with caution.

Eccentric exercises

Four studies evaluated the outcome of eccentric exercises for insertional Achilles tendinopathy [13, 22, 26, 42]. There were some important differences between treatment protocols; most notably, the difference between full range

of motion exercises and floor level exercises. Evaluating the separate results an important difference was found: good outcomes for floor level exercises compared to insufficient results of full range of motion studies (Table 5). All studies used a 12-week training protocol with painful exercises 2 or 3 times daily, with up to three sets of 10–15 repetitions each. However, all studies had a relative short follow-up as the longest was 4 months. Despite the good outcome for midportion Achilles tendinopathy with full range of motion eccentric exercises, and in contrast with the statement of a previous systematic review, full range eccentric training may not be the first treatment of choice for insertional Achilles tendinopathy as the patient satisfaction is unfavourable compared to other conservative treatment modalities for insertional Achilles tendinopathy [24]. 70 % of all patients were unsatisfied after a 12-week full range eccentric training regime (Table 5). Although evaluated in only one study, the results of floor level eccentric exercises are much more promising: 67 % per cent satisfied after floor level compared to 30 % satisfied after full range exercises. The focus of eccentric treatment should therefore be on floor level exercises.

Extracorporeal shockwave therapy

One prospective study and one randomized controlled trial evaluated ESWT for insertional Achilles tendinopathy [14, 42]. The same study group performed both studies. A low-to-medium energy ESWT was given without the use of anaesthesia. The RCT and prospective study both showed good results. A decrease in subjective pain scale (−4.8) and high patient satisfaction were found (Table 5). For the prospective study, it should be noted that the patient selection was biased on insurance claim and the control group received not more than a wait-and-see protocol, the outcome may therefore be affected by placebo effect.

Others

One study evaluated Laser CO² with TECAR and cryoultrasound [10]. All showed significant decrease on a VAS. Although the results seem promising, the study included only 15 patients with insertional problems, the three different groups consisted of merely five patients. No other studies evaluated the outcome of these treatments; additional evidence is a necessity as the current evidence is too weak to draw conclusions or to provide a clinical treatment advice.

Calcifications versus no calcifications

Nine of 14 studies reported on the presence of a bony disorder, located at the insertion of the AT into the

calcaneus, in every case [12, 15, 20, 22, 26, 31, 37, 42, 53]. Four of six studies evaluating surgery for insertional Achilles tendinopathy reported on bony disorders [12, 20, 30, 53]. In addition, the diagnostic method was based on imaging in 12 studies, and this was unclear in two other studies (Table 3). This is important, as the outcome of a treatment could be different in patients *with* bony pathology compared to a group *without* bony pathology. Hence, imaging is essential in the diagnoses and thus treatment for insertional Achilles tendinopathy. Future research should differentiate between and/or compare treatments in a group *with* bony disorder and a group *without* bony disorder.

Study limitations and clinical relevance

The most important limitation of this study is the low level of evidence of included studies and the substantial risk of bias herein. Only one randomized controlled trial could be included; most studies were prospective or retrospective in design. Another important limitation is the terminological confusion surrounding Achilles tendon pathology [18, 19, 30, 38, 50]. This systematic review used strict inclusion criteria to define insertional Achilles tendinopathy; as a result, studies may be excluded based on the used terminology. As insertional Achilles tendinopathy differs from other chronic Achilles tendon pathology, the results of this analysis are largely dependent on the proper diagnosis and terminology of the included studies [30, 31, 35, 45, 50, 52]. Although all studies reported that the diagnosis was based on clinical findings and different imaging techniques, patient' symptoms may have been based on other pathology instead of insertional Achilles tendinopathy (e.g. retrocalcaneal bursitis). Finally, insertional Achilles tendinopathy may occur together with retrocalcaneal bursitis, if patients have this combination and are treated for one or the other a minimal effect of intervention can be expected [1]. A comparison between these entities after surgical intervention was made by Watson [52]; he found different results between these pathologies in spite of the same surgical intervention, hereby reemphasizing the difference between these pathologies.

Despite the generally low level of quality of available studies, the outcome is of clinical importance [4]. This study is the first to evaluate the effectiveness of non-surgical and surgical interventions for insertional Achilles tendinopathy based on patient satisfaction, pain and complication rates. In addition, it points at the clear distinction (in outcome) between the currently used eccentric exercises protocols for insertional Achilles tendinopathy; finally, it shows the importance of differentiation between calcified and non-calcified insertional Achilles tendinopathy.

Conclusion

It was found that there are numerous treatment modalities, non-surgical and surgical, for insertional Achilles tendinopathy. Regarding non-surgical treatment, although both showed a decrease in VAS score, a substantial difference was found between the outcome of full range eccentric exercises and floor level exercises in insertional Achilles tendinopathy; full range of motion exercises resulted in a low patient satisfaction compared to floor level exercises or other treatment modalities. The results of floor level eccentric exercises are promising but currently evaluated by only one study. ESWT, although based on only two studies, appears effective in patients with non-calcified insertional Achilles tendinopathy. Evaluation by other study groups would substantially strengthen the current positive position of ESWT. Different surgical techniques are used to treat insertional Achilles tendinopathy. There are some differences in outcome, mainly regarding complication rate. Despite these differences, the patient satisfaction is relatively high in all surgical studies. Based on the available studies, it is not possible to draw any conclusion regarding the best surgical technique for insertional Achilles tendinopathy. Finally, it is evident that more high level evidence studies are necessary to be more conclusive regarding the best available treatment. Future studies should focus on clearly stating the exact studied pathology (bony involvement/not) and use validated outcome measures.

Conflict of interest No financial or material support was received.

References

1. Aaron DL, Patel A, Kayiaros S, Calfee R (2011) Four common types of bursitis: diagnosis and management. *J Am Acad Orthop Surg* 19:359–367
2. Aronow MS (2005) Posterior heel pain (retrocalcaneal bursitis, insertional and noninsertional Achilles tendinopathy). *Clin Podiatr Med Surg* 22:19–43
3. Astrom M, Westlin N (1992) No effect of piroxicam on achilles tendinopathy. A randomized study of 70 patients. *Acta Orthop Scand* 63:631–634
4. Bhandari M, Montori VM, Devereaux PJ, Wilczynski NL, Morgan D, Haynes RB (2004) Doubling the impact: publication of systematic review articles in orthopaedic journals. *J Bone Joint Surg Am* 86-A:1012–1016
5. Calder JD, Saxby TS (2003) Surgical treatment of insertional Achilles tendinosis. *Foot Ankle Int* 24:119–121
6. Carmont MR, Maffulli N (2007) Management of insertional Achilles tendinopathy through a Cincinnati incision. *BMC Musculoskelet Disord* 8:82
7. Chiara VM, Guzzini M, Ferretti A (2003) Operative treatment of chronic Achilles tendinopathy. *Int Orthop* 27:307–310
8. Clain MR, Baxter DE (1992) Achilles tendinitis. *Foot Ankle* 13:482–487
9. Coleman BD, Khan KM, Maffulli N, Cook JL, Wark JD (2000) Studies of surgical outcome after patellar tendinopathy: clinical

- significance of methodological deficiencies and guidelines for future studies. Victorian Institute of Sport Tendon Study Group. *Scand J Med Sci Sports* 10:2–11
10. Costantino C, Pogliacomì F, Vaianti E (2005) Cryoultrasound therapy and tendonitis in athletes: a comparative evaluation versus laser CO₂ and t.e.ca.r. therapy. *Acta Biomed* 76:37–41
 11. de Vos RJ, Weir A, van Schie HT, Bierma-Zeinstra SM, Verhaar JA, Weinans H, Tol JL (2010) Platelet-rich plasma injection for chronic Achilles tendinopathy: a randomized controlled trial. *JAMA* 303:144–149
 12. Elias I, Raikin SM, Besser MP, Nazarian LN (2009) Outcomes of chronic insertional Achilles tendinosis using FHL autograft through single incision. *Foot Ankle Int* 30:197–204
 13. Fahlstrom M, Jonsson P, Lorentzon R, Alfredson H (2003) Chronic Achilles tendon pain treated with eccentric calf-muscle training. *Knee Surg Sports Traumatol Arthrosc* 11:327–333
 14. Furia JP (2005) Extracorporeal shockwave therapy in the treatment of chronic insertional Achilles tendinopathy. *Orthopade* 34:571–578
 15. Furia JP (2006) High-energy extracorporeal shock wave therapy as a treatment for insertional Achilles tendinopathy. *Am J Sports Med* 34:733–740
 16. Furia JP (2008) High-energy extracorporeal shock wave therapy as a treatment for chronic noninsertional Achilles tendinopathy. *Am J Sports Med* 36:502–508
 17. Gerken AP, McGarvey W, Baxter DE (1996) Insertional Achilles tendinitis. *Foot Ankle Clin* 1:237–248
 18. Harris CA, Peduto AJ (2006) Achilles tendon imaging. *Australas Radiol* 50:513–525
 19. Jerosch J, Schunck J, Sokkar SH (2007) Endoscopic calcaneoplasty (ECP) as a surgical treatment of Haglund's syndrome. *Knee Surg Sports Traumatol Arthrosc* 15:927–934
 20. Johnson KW, Zalavras C, Thordarson DB (2006) Surgical management of insertional calcific achilles tendinosis with a central tendon splitting approach. *Foot Ankle Int* 27:245–250
 21. Jones DC, James SL (1984) Partial calcaneal osteotomy for retrocalcaneal bursitis. *Am J Sports Med* 12:72–73
 22. Jonsson P, Alfredson H, Sunding K, Fahlstrom M, Cook J (2008) New regimen for eccentric calf-muscle training in patients with chronic insertional Achilles tendinopathy: results of a pilot study. *Br J Sports Med* 42:746–749
 23. Kane TP, Ismail M, Calder JD (2008) Topical glyceryl trinitrate and noninsertional Achilles tendinopathy: a clinical and cellular investigation. *Am J Sports Med* 36:1160–1163
 24. Kearney R, Costa ML (2010) Insertional achilles tendinopathy management: a systematic review. *Foot Ankle Int* 31:689–694
 25. Kirkup J (1988) Mythology and History. In: Helal B, Wilson D (eds) *The foot*. Churchill Livingstone, Philadelphia, pp 1–8
 26. Knobloch K (2007) Eccentric training in Achilles tendinopathy: is it harmful to tendon microcirculation? *Br J Sports Med* 41:e2
 27. Knobloch K, Grasemann R, Spies M, Vogt PM (2008) Midportion achilles tendon microcirculation after intermittent combined cryotherapy and compression compared with cryotherapy alone: a randomized trial. *Am J Sports Med* 36:2128–2138
 28. Leach RE, Dilorio E, Harney RA (1983) Pathologic hindfoot conditions in the athlete. *Clin Orthop Relat Res* 177:116–121
 29. Leitze Z, Sella EJ, Aversa JM (2003) Endoscopic decompression of the retrocalcaneal space. *J Bone Joint Surg Am* 85:1488–1496
 30. Maffulli N, Khan KM, Puddu G (1998) Overuse tendon conditions: time to change a confusing terminology. *Arthroscopy* 14:840–843
 31. Maffulli N, Testa V, Capasso G, Sullo A (2004) Calcific insertional Achilles tendinopathy: reattachment with bone anchors. *Am J Sports Med* 32:174–182
 32. McGarvey WC, Palumbo RC, Baxter DE, Leibman BD (2002) Insertional Achilles tendinosis: surgical treatment through a central tendon splitting approach. *Foot Ankle Int* 23:19–25
 33. McLauchlan GJ, Handoll HH (2001) Interventions for treating acute and chronic Achilles tendinitis. *Cochrane Database Syst Rev* 10(8):CD000232
 34. Morelli V, James E (2004) Achilles tendonopathy and tendon rupture: conservative versus surgical management. *Prim Care* 31:1039–1054
 35. Myerson MS, McGarvey W (1999) Disorders of the Achilles tendon insertion and Achilles tendinitis. *Instr Course Lect* 48:211–218
 36. Norregaard J, Larsen CC, Bieler T, Langberg H (2007) Eccentric exercise in treatment of Achilles tendinopathy. *Scand J Med Sci Sports* 17:133–138
 37. Ohberg L, Alfredson H (2003) Sclerosing therapy in chronic Achilles tendon insertional pain—results of a pilot study. *Knee Surg Sports Traumatol Arthrosc* 11:339–343
 38. Paavola M, Jarvinen TA (2005) Paratendinopathy. *Foot Ankle Clin* 10:279–292
 39. Petersen W, Welp R, Rosenbaum D (2007) Chronic Achilles tendinopathy: a prospective randomized study comparing the therapeutic effect of eccentric training, the AirHeel brace, and a combination of both. *Am J Sports Med* 35:1659–1667
 40. Philippot R, Wegrzyn J, Grosclaude S, Besse JL (2010) Repair of insertional achilles tendinosis with a bone-quadriceps tendon graft. *Foot Ankle Int* 31:802–806
 41. Puddu G, Ippolito E, Postacchini F (1976) A classification of Achilles tendon disease. *Am J Sports Med* 4:145–150
 42. Rompe JD, Furia J, Maffulli N (2008) Eccentric loading compared with shock wave treatment for chronic insertional achilles tendinopathy. A randomized, controlled trial. *J Bone Joint Surg Am* 90:52–61
 43. Rompe JD, Furia J, Maffulli N (2009) Eccentric loading versus eccentric loading plus shock-wave treatment for midportion achilles tendinopathy: a randomized controlled trial. *Am J Sports Med* 37:463–470
 44. Ryan M, Wong A, Taunton J (2010) Favorable outcomes after sonographically guided intratendinous injection of hyperosmolar dextrose for chronic insertional and midportion achilles tendinosis. *AJR Am J Roentgenol* 194:1047–1053
 45. Schepsis AA, Jones H, Haas AL (2002) Achilles tendon disorders in athletes. *Am J Sports Med* 30:287–305
 46. Stergioulas A, Stergioulas M, Aarskog R, Lopes-Martins RA, Bjordal JM (2008) Effects of low-level laser therapy and eccentric exercises in the treatment of recreational athletes with chronic achilles tendinopathy. *Am J Sports Med* 36:881–887
 47. Tallon C, Coleman BD, Khan KM, Maffulli N (2001) Outcome of surgery for chronic Achilles tendinopathy. A critical review. *Am J Sports Med* 29:315–320
 48. Thomas JL, Christensen JC, Kravitz SR, Mendicino RW, Schuberth JM, Vanore JV, Weil LS Sr, Zlotoff HJ, Bouche R, Baker J (2010) The diagnosis and treatment of heel pain: a clinical practice guideline—revision 2010. *J Foot Ankle Surg* 49:S1–S19
 49. van Dijk CN, van Dyk GE, Scholten PE, Kort NP (2001) Endoscopic calcaneoplasty. *Am J Sports Med* 29:185–189
 50. van Dijk CN, van Sterkenburg MN, Wiegerinck JJ, Karlsson J, Maffulli N (2011) Terminology for Achilles tendon related disorders. *Knee Surg Sports Traumatol Arthrosc* 19(5):835–841
 51. Wagner E, Gould JS, Kneidel M, Fleisig GS, Fowler R (2006) Technique and results of Achilles tendon detachment and reconstruction for insertional Achilles tendinosis. *Foot Ankle Int* 27:677–684
 52. Watson AD, Anderson RB, Davis WH (2000) Comparison of results of retrocalcaneal decompression for retrocalcaneal bursitis and insertional achilles tendinosis with calcific spur. *Foot Ankle Int* 21:638–642
 53. Yodlowski ML, Scheller AD Jr, Minos L (2002) Surgical treatment of Achilles tendinitis by decompression of the retrocalcaneal bursa and the superior calcaneal tuberosity. *Am J Sports Med* 30:318–321